

### **AMENDMENTS TO THE CLAIMS:**

This listing of claims will replace all prior versions, and listings, of claims in the application:

#### **Listing of the claims:**

1. (Currently amended) A method for screening compounds useful for the treatment of proliferative and differentiative disorders comprising contacting a compound with a cell or a cell extract expressing either Cks1 and Skp2, or Cks1, p27 and Skp2, and detecting a change in the activity of Skp2.
2. (Original) The method of Claim 1 wherein the change in the activity of Skp2 is detected by detecting a change in the interaction of Skp2 with either p27 or Cks1.
3. (Original) The method of Claim 1 wherein the change in the activity of Skp2 is detected by detecting a change in the ubiquitination of p27 or degradation of p27 or Cks1.
4. (Currently amended) A method for screening compounds useful for the treatment of proliferative and differentiative disorders comprising adding a compound in a purified system containing either Cks1 and Skp2, or Cks1, p27 and Skp2, and detecting a change in the activity of Skp2.
5. (Original) The method of Claim 4 wherein the change in the activity of Skp2 is detected by detecting a change in the interaction of Skp2 with either p27 or Cks1.
6. (Original) The method of Claim 4 wherein the change in the activity of Skp2 is detected by detecting a change in the ubiquitination of p27 or degradation of p27 or Cks1.
7. (Currently amended) A method for screening compounds useful for the treatment of proliferative and differentiative disorders comprising:
  - (a) adding a compound in a purified system containing Skp2 and one or both of: (i) a polypeptide corresponding to the carboxy terminus of the human p27 chain having the sequence NAGSVEWTPKKPGLRRRQT (SEQ. ID. NO: 91) with or without a phosphothreonine at position 187 8 and (ii) Cks1; and

(b) detecting a change in the activity of Skp2.

8. (Original) The method of Claim 7 wherein the change in the activity of Skp2 is detected by detecting a change in the interaction of Skp2 with either the polypeptide or Cks1.

9. (Original) The method of Claim 7 wherein the change in the activity of Skp2 is detected by detecting a change in the ubiquitination of the polypeptide or degradation of the polypeptide or Cks1.